

KETO E. MSHIGENI & REGINE JAHN

***Eucheuma* (Rhodophyta, Gigartinales) in the W Indian Ocean region - notes on collections kept in Berlin-Dahlem and Hamburg**

**Abstract**

Mshigeni, K. E. & Jahn, R.: *Eucheuma* (Rhodophyta, Gigartinales) in the W Indian Ocean region - notes on collections kept in Berlin-Dahlem and Hamburg. - Willdenowia 25: 399-407. 1995. - ISSN 0511-9618.

A survey on the *Eucheuma* collections from the W Indian Ocean region kept at B and HBG is given; notes on the distribution of *E. denticulatum*, *E. odontophorum* and *E. platycladum* are added and the characters distinguishing *E. odontophorum* from *E. platycladum* listed.

**Introduction**

The red algal genus *Eucheuma* J. Agardh (Gigartinales, Solieriaceae) is economically the most important seaweed in the western Indian Ocean region. Some of its species are exported by Tanzania (Anderson 1953; Mshigeni 1973, 1987) and Madagascar (Mollion, Andriantsiferana & Sekkal 1990; Rabesandratana & Rabesandratana 1992) for use as a source of carrageenan (Doty 1973, 1988; Mshigeni & Semesi 1977). The commercial importance of *Eucheuma* in the region is likely to increase in the near future with the introduction and successful assimilation of its cultivation technology in Tanzania's rural communities along the coast (Mshigeni 1992). This *Eucheuma* farming technology has been pioneered by M. Doty and V. Alvarez (Doty 1973) in the Indo-Pacific region.

However, the various species of *Eucheuma* in the W Indian Ocean region are still imperfectly known (Doty 1988; Mollion, Andriantsiferana & Sekkal 1990), and there is a need for a detailed study of their taxonomy and distribution. Most of the earlier authors, who validated the names of *Eucheuma* species from the region, based their descriptions on dead specimens or just fragments. Often the samples were collected by non-specialists or short-term visitors, who did not record field data and the morphological variability of the sampled plants in their natural habitats. Therefore, they could not know to what extent the morphological characters had been influenced by the environment (e.g. exposure to waves, intertidal stress, grazing) or by age phenomena. The current circumscription of the *Eucheuma* species of this region (Mshigeni 1984, 1987) were reached after extensive field studies of these taxa in East Africa and Mauritius, and by comparing the collections with specimens from other herbaria (listed also in Doty 1988).

The present study focusses on the historical *Eucheuma* material from the W Indian Ocean region in the herbaria of Berlin-Dahlem (B) and Hamburg (HBG).

**Results and discussion**

Our results are summarized in two tables. Tab. 1 contains the details of all *Eucheuma* specimens of African origin that were found in the herbaria investigated. Tab. 2 lists published records of distribution of if cut at a distance and axiferous when cut close to the apex. He has revised the sectional arrangement of the various species of *Eucheuma* and separated some of the taxa into the new genus *Kappaphycus* Doty, which encompasses species characterised by the presence of kappa carrageenan (e.g. *K. cottonii*, *K. inermis*). But he does not place *Eucheuma platycladum* under *Kappaphycus*, even though the species produces kappa carrageenan preponderantly (see also Doty & Santos 1978; Mshigeni & Semesi 1977, Doty 1988: 204). Since the precise chemical analysis of the carrageenan types in the majority of the *Eucheuma* species reported in the region is still to be done, and since some findings are not adequately conclusive, we have retained the old genus name.

***Eucheuma denticulatum* (Burman) Collins & Hervey**

The locality of the two specimens kept at B, i.e. Cape of Good Hope or just Cap., is doubtful, since the southern tip of Africa is believed to be too cold for *Eucheuma*. As shown in Tab. 2, and pointed out by Mshigeni (1979, 1984, 1987) and Jaasund (1976, 1977), the species is widely distributed in East Africa. Its range has recently been extended to Madagascar (Mollion, Andriantsiferana *Eucheuma* species from the western Indian Ocean region; in addition to the cited species, Doty (1988) recognises the occurrence of *Eucheuma serra* J. Agardh in Mauritius.

The various species of *Eucheuma* were arranged by Weber-van Bosse (1928) into two sections: section *Axifera* encompasses species with a central axis of relatively small and densely packed cells (e.g. *E. denticulatum*, *E. serra*, *E. horridum*); section *Anaxifera* embraces taxa (e.g. *E. cottonii*, *E. platycladum*) whose fronds lack a definite central axis. But Doty (1988: 204-205) has cautioned that transverse sections of the same thallus may appear anaxiferous & Sekkal 1990; Rabesandratana & Rabesandratana 1992). In addition we found a box with dried specimens of *E. denticulatum* which lacked a label; this might be some of the material which Schmitz (1895: 139) referred to as *E. spinosum*.

### *E. odontophorum* Børgesen

The record from Lamu in Kenya (Fig. 1) represents the northernmost locality of this species. Doty (1988) has also documented its occurrence in Mauritius but writes: "... Not enough is known about *E. odontophorum* to circumscribe it in detail. The type specimen ... and the only other collection seen ... is of thin axiferous, strap-shaped fragmentary blades, ornamented only by lateral teeth, so as to look like a juvenile form of *E. platycladum*". He points out that the placement of *E. platycladum* in the sections is unsatisfactory, since there are mixed reports of their axes being anaxiferous as well as axiferous. Earlier Weber-van Bosse (1928: 417) was not sure of the placement of this species in the section *Anaxifera* since she was unable to locate the type specimen in Berlin.

### *E. platycladum* Schmitz

In our search for *Eucheuma* specimens in Berlin, we detected a small cardboard box containing a dried specimen (Fig. 2A) with spinulose and compressed, anastomosed axes, which perfectly fitted our conception of *E. platycladum* Schmitz. The handwritten label bears the collectors name, the number and type locality indicated by Schmitz (1895: 152) for *E. platycladum*: Holst 1340, Dar es Salaam. But the species is labelled *Eucheuma spinosum* (L.) J. Agardh as Schmitz (1895: 152) explains: "This *Eucheuma* species is quite a big plant of peculiar habit. I have already seen several specimens of it, and I am astonished that nobody has described it as a separate species. I found it [the type specimen] named *E. spinosum*, but it has no similarity to this species; it rather resembles the Southwest-Australian *E. speciosum* (Sond.) J. Ag., but differs also slightly from this species". Since Schmitz died unexpectedly (see footnote in Schmitz 1895: 137), the label on the specimen was apparently not changed. The cardboard box as well as the printed label with the locality "D.[eutsch] Ostafrika" (German East Africa) indicates that this specimen was used as exhibition material, some of which survived the 1943 bombing of the Botanical Museum Berlin-Dahlem. This would also explain why Weber-van Bosse (1928: 417) was unable to locate the type. If duplicates mounted on herbarium sheets with correct labels had existed, she would have seen them; but they would have been destroyed together with those of *E. inerme*. What a coincidence that the re-discovery date, December 1992, was exactly a century after Holst had collected the material in December 1892 (Schmitz 1895: 138)!

Besides material used for exhibition purposes in the Botanical Museum Berlin-Dahlem before the Second World War most of the wet collections remained extant; among them we found another specimen of *E. platycladum* collected by A. Voeltzkow in 1903 from the "Tamatave Riff", Madagascar (Fig. 2B, Tab. 1) which confirms the recent report by Mollion, Andriantsiferana & Sekkal (1990) on the existence of *E. platycladum* in Madagascar, thus extending its geographical range beyond East Africa.

Not a single specimen of this species was found pressed onto herbarium sheets. This might be due partly to the inherent difficulties with drying its thalli, because of the cartilaginous nature of its fronds, the anastomosing habit, and the high phycocolloid content. The coastal inhabitants in Tanzania refer to the seaweed as mwani-mkate, meaning bread-like seaweed, or mwani-kisahani, referring to its plate-like appearance. Fronds of the alga form expansive cake-like masses over the substrata, sometimes forming cushions which weigh several kilogrammes.

We do not share the view that the morphological differences of *E. odontophorum* and *E. platycladum* are not clear-cut (Doty 1988: 198). Both species have dorsiventrally compressed fronds and are spinulose, but there are differences: the axes of *E. odontophorum* are axiferous, those of *E. platycladum* anaxiferous; the fronds of *E. odontophorum* show distinct segmentation (Fig. 1) those of *E. platycladum* lack constrictions along the axes; the younger axes of *E. odontophorum* are almost cylindrical and the older sub-cylindrical, the axes of *E. platycladum* exhibit a stronger tendency of flattening and anastomosing, even on the relatively young branchlets; the axes of *E. odontophorum* are terminally downwardly arched (Mshigeni 1979, fig. 11; Mshigeni & Semesi 1977, fig. 5), this is not so distinct in the fronds of *E. platycladum*.

### Acknowledgements

Thanks are due to the Institut für Allgemeine Botanik, University of Hamburg, for the use of the research facilities; to Dr. B. Zimmer for useful suggestions, and to B. Schreiber for assisting with the photographic work. The senior author is most grateful to the German Academic Exchange Service (DAAD) for financial support.

### References

- Agardh, J. G. 1892: *Analecta algologica* 1. - Lund.
- Anderson, G. W. 1953: A note on the seaweed resources of Zanzibar Protectorate. - *Proc. Int. Seaweed Symp.* 1: 102-103.
- Børgesen, F. 1943: Some marine algae of Mauritius. III. *Rhodophyceae*, Part. 2. *Gelidiales*, *Cryptonemiales*, *Gigartinales*. - *Biol. Meddel. Kongel. Danske Vidensk. Selsk.* 19(1).
- 1950: Some marine algae of Mauritius. Additions II. - *Biol. Meddel. Kongel. Danske Vidensk. Selsk.* 18(11).

- Dickie, G. 1875: On the algae of Mauritius. - J. Linn. Soc., Bot. 1: 190-202.
- Dixon, P. S. 1962: Taxonomic and nomenclatural notes on the *Florideae* III. - Bot. Not. 115: 245-260.
- Doty, M. S. 1973: Farming the red seaweed *Eucheuma* for carrageenans. - Micronesica 9: 59-73.
- 1988: Prodromus ad systematica Eucheumatoiderum: a tribe of commercial seaweeds related to *Eucheuma* (*Soleriaceae*, *Gigartinales*), p. 159-207. - In: Abbott, I. A. (ed.), Taxonomy of economic seaweeds, with special reference to some Pacific and Caribbean species 2. - La Jolla, Calif.
- & Santos, G. A. 1978: Carrageenans from tetrasporic and cystocarpic *Eucheuma* species. - Aquatic Bot. 4: 143-149.
- Harvey, W. H. 1834: Notice on a collection of algae communicated to Dr. Hooker by the late Mrs. Charles Telfair from Cap Malheureux in Mauritius. - J. Bot. (Hooker) 1: 147-157.
- Jaasund, E. 1976: Intertidal seaweeds in Tanzania: a field guide. - Tromsø.
- 1977: Marine algae in Tanzania VII. - Bot. Mar. 20: 415-425.
- Mollion, J., Andriantsiferana, M. & Sekkal, M. 1990: A study of the phycocolloids from *Gelidium madagascariense* and *Eucheuma denticulatum* (*Rhodophyta*) collected on the south coasts of Madagascar. - Hydrobiologia 204/205: 655-659.
- Mshigeni, K.E. 1973: Exploitation of seaweeds in Tanzania: the genus *Eucheuma* and other algae. - Tanzania Notes Rec. 72: 19-36.
- 1979: The economic algal genus *Eucheuma* (*Rhodophyta*, *Gigartinales*): observations on Tanzanian species. - Bot. Mar. 22: 437-445.
- 1984: The red algal genus *Eucheuma* (*Rhodophyta*, *Gigartinales*) in East Africa: an under-exploited resource. - Proc. Int. Seaweed Symp. 11: 347-350.
- 1987: The taxonomy, ecology, and economic potential of the red seaweed *Eucheuma* J. Agardh in the western Indian Ocean region. - Techn. Publ. Ser. Commonw. Sci. Council 221: 1-82.
- 1992: Seaweed farming in Tanzania: a success story, p. 221-245. - In: Mshigeni, K. E. et al. (eds.): Proceedings of the First International Workshop on Sustainable Seaweed Resource Development in Sub-Saharan Africa, Windhoek, Namibia. - [s.l.].
- & Semesi, A. K. 1977: Studies on carrageenans from the economic red algal genus *Eucheuma* in Tanzania. - Bot. Mar. 20: 239-242.
- Rabesandratana, H. D. & Rabesandratana, R. N. 1992: Seaweeds and their uses in Madagascar, p. 297-307. - In: Mshigeni, K. E. et al. (eds.), Proceedings of the First International Workshop on Sustainable Seaweed Resource Development in Sub-Saharan Africa, Windhoek, Namibia. - [s.l.].
- Schmitz, F. 1895: Marine Florideen von Deutsch Ost-Afrika. - Bot. Jahrb. 21: 137-177.
- Sonder, W. 1879: *Algae*, p. 79-85. - In: Kersten, O., Baron Carl Claus von den Deckens Reisen in Ost-Afrika in den Jahren 1859-1865 3(3). - Leipzig.
- Weber-van Bosse, A. 1913: Marine algae, *Rhodophyceae* of the Sealark-Expedition collected by Mr. J. Stanley Gardiner, M.A. - Trans. Linn. Soc. London, ser. 2, Bot. 8: 105-142.
- 1928: Liste des algues du Siboga. IV. Part 3. *Gigartinales* et *Rhodymeniales*, p. 393-533. - In: Weber, M. (ed.), Siboga Exped. Monogr. 59d.
- Yarish, C. & Wamukoya, G. 1990: Seaweeds of potential economic importance in Kenya: field survey and future prospects. - Hydrobiologia 204/205: 339-346.

Addresses of the authors:

K.E. Mshigeni, Department of Botany, University of Namibia, Private Bag 13301, Windhoek, Namibia.

R. Jahn, Botanischer Garten und Botanisches Museum Berlin-Dahlem, Freie Universität Berlin, Königin-Luise-Str. 6-8, D-14191 Berlin.

Tab. 1: *Eucheuma* specimens from the W Indian Ocean region in B and HBG.

**Locality//Collector//Date//Voucher//Remarks**

***Eucheuma denticulatum***

Cape of Good Hope//[Herb. Schiffner]//Not recorded//B: Algae 15031//Non-compressed, spinulose, verticillate

Cape of Good Hope//Not recorded//Not recorded//B: Algae 23778//Non-compressed, spinulose, verticillate

***E. horridum***

Mauritius//not recorded//not recorded//B: Algae 15042// Spinulose, non-compressed, some of the spines elongated into branchlets, irregularly arranged

Madagascar//herb. Wollny; herb. Th. Reinbold//Not recorded//HBG: H.H.10//Fronds slender, spinulose, non-compressed, spines not arranged in regular whorls

***E. odontophorum***

Kiu Island Lagoon, Kenya// P.J. Greenway 9388 & S.P. Rawlins//13.10.1957//B: Algae 33866, 34190//Fronds clad with spines, semi-compressed, with many constrictions along the axes

***E. platycladum***

Dar es Salaam, Tanzania// Holst 1340//12.1892//B: Algae 36507//Type; single specimen, spinulose, axes dorsiventrally compressed, constrictions along the axes not distinct

Tamatave Riff, Madagascar//Voeltzkow//1903//B: Algae 37362 (wet)//Fronds clad with spines, dorsiventrally compressed, constrictions along thallus axes not distinct

***E. striatum***

Zanzibar//Stuhlmann//1889?//HBG: H.H. 21//Type (fragment)

Pemba Island//Voeltzkow//1903//B: Algae 29 068//Several specimens, non-spinulose, non-compressed, with acuminate branches

Tab. 2: Published records of *Euclidean* species in the W Indian Ocean region (see Dixon (1962) and Mshigeni (1987) for synonymy and re-identifications), \*= records need confirmation.

**Locality//Reference//Name recorded//Collector//Date**

***Euclidean cottonii***

Mascarene Ridge//Weber-van Bosse (1913)//*E. cottonii*//J.S. Gardiner//Sealark-Expedition

Tanzania//Mshigeni (1973; 1979)//*E. okamurai*//K.E. Mshigeni//1969

Tanzania//Mshigeni & Semesi (1977)//*E. okamurai*//K.E. Mshigeni//1969

Tanzania//Jaasund (1976, 1977)//*E. okamurai*//E. Jaasund//1966-69

Kenya\*/Yarish & Wamukoya (1990)//*E. cottonii*

***E. denticulatum***

Mauritius//Dickie (1875)//*E. spinosum*//Pike//1869 ?

Zanzibar//Sonder (1879)//*E. spinosum*//A. Roscher//1859

Zanzibar//Schmitz (1895)//*E. spinosum*//?1885

Tanzania//Schmitz (1895)//*E. spinosum*//C. Holst//1892

Tanzania//Mshigeni (1973, 1979)//*E. spinosum*//K.E. Mshigeni//1969

Tanzania//Mshigeni & Semesi (1977)//*E. spinosum*//K.E. Mshigeni//1969

Tanzania//Jaasund (1976), (1977)//*E. spinosum*//E. Jaasund//1966-69

***E. horridum***

Mauritius//Harvey (1834)//*Sphaerococcus horrida* //Ch. Telfair//18??

Mauritius//Dickie (1875)//*E. horridum*//Pike//1869?

Mauritius//J. Agardh (1892)//*E. jugatum*

Mauritius//Børgesen (1943)//*E. serra*//Th. Mortensen//1929

Mauritius//Børgesen (1943, 1950)//*E. jugatum* //Th. Mortensen//1929

Mauritius//Børgesen (1943, 1950)//*E. horridum*//Th. Mortensen//1929

Tanzania//Mshigeni (1987)//*E. horridum*//K.E. Mshigeni//1969

Kenya//Yarish & Wamukoya (1990)//*E. horridum*

Madagascar\*/Mollion, Andriantsiferana & Sekkal (1990)//*E. denticulatum*

Madagascar\*/Rabesandratana & Rabesandratana (1992)//*E. denticulatum*

***E. odontophorum***

Mauritius//Dickie (1875)//*E. spinosum*//Pike//1869?

Mauritius//Børgesen (1943)//*E. odontophorum*//Th. Mortensen//1929

Mauritius//Børgesen (1943)//*E. speciosum* (Sonder) var. *mauritiana*//Th. Mortensen //1929

Tanzania//Mshigeni (1973; 1979)//*E. speciosum* var. *mauritiana*//K.E. Mshigeni//1969

Tanzania//Mshigeni & Semesi (1977)//*E. speciosum* var. *mauritiana*//K.E. Mshigeni//1969

Tanzania//Jaasund (1976, 1977)//*E. speciosum* var. *mauritiana*//*E. Jaasund*//1966-69  
Tanzania//Mshigeni (1987)//*E. odontophorum*//*K.E. Mshigeni*//1969  
Mauritius//Doty (1988)//*E. odontophorum*//*M. Doty*

***Eucheuma platycladum***

Tanzania//Schmitz (1895)//*E. platycladum*//*C. Holst*//10.1892  
Tanzania//Mshigeni (1973; 1979)// *E. platycladum* //*K.E. Mshigeni*//1969  
Tanzania//Mshigeni & Semesi (1977)//*E. platycladum*//*K.E. Mshigeni*//1969  
Tanzania//Jaasund (1976, 1977)//*E. platycladum*//*E. Jaasund*//1966-69  
Tanzania//Mshigeni (1987)//*E. platycladum*//*K.E. Mshigeni*//1969  
Tanzania//Doty (1988)//*E. platycladum*//*G.F.Papenfuss & R.F. Scagel*//1962  
Tanzania//Doty (1988)//*E. platycladum*//*K.E. Mshigeni*//1969  
Madagascar//Mollion, Andriantsiferana & Sekkal (1990)//*E. platycladum*  
Madagascar//Rabesandratana & Rabesandratana (1992)//*E. platycladum*

***E. striatum***

Zanzibar//Schmitz (1895)//*E. inerme*////1885  
Tanzania//Schmitz (1895)//*E. inerme*//*C. Holst*//1885  
Zanzibar//Schmitz (1895)//*E. striatum*//*F. Stuhlmann*//1888-90  
Tanzania//Mshigeni (1973, 1979)//*E. striatum*//*K.E. Mshigeni*//1969  
Tanzania//Mshigeni & Semesi (1977)//*E. striatum*//*K.E. Mshigeni*//1969  
Tanzania//Jaasund (1976, 1977)//*E. striatum*//*E. Jaasund*//1966-69  
Madagascar//Mollion, Andriantsiferana & Sekkal (1990)//*E. striatum*  
Madagascar//Rabesandratana & Rabesandratana (1992)//*E. striatum*